



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11D 3/386, C12N 9/24, 9/26	A1	(11) International Publication Number: WO 96/13568 (43) International Publication Date: 9 May 1996 (09.05.96)
(21) International Application Number: PCT/US95/12490 (22) International Filing Date: 29 September 1995 (29.09.95) (30) Priority Data: 94870169.3 27 October 1994 (27.10.94) EP (34) Countries for which the regional or international application was filed: BE et al. (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BAECK, Andre, Cesar [BE/BE]; Putsesteenweg 273, B-2820 Bonheiden (BE). BUSCH, Alfred [DE/BE]; Handelsstraat 210, B-1840 Londerzeel (BE). VERSCHUERE, Ann, Katrien, Marie, Agnes [BE/BE]; Reigerlo 19, B-8730 Beernem (BE). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: BR, CA, CN, JP, MX, US, VN. Published <i>With international search report.</i>
(54) Title: CLEANING COMPOSITIONS COMPRISING XYLANASES (57) Abstract The invention concerns cleaning compositions comprising enzymes having xylanolytic activity. More in detail the invention relates to detergent compositions preferably compact detergents having xylanase enzyme incorporated. Said compositions show an excellent boost on cleaning performance on fruit, vegetables and/or mud and clay compounds containing said soils.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

CLEANING COMPOSITIONS COMPRISING XYLANASES

Field of the Invention

The present invention relates to cleaning compositions, including dishwashing, laundry compositions and oral/dental compositions as well, containing an enzyme with xylanolytic activity.

More in particular, the invention relates to compact detergent compositions, granules and liquids as well having said enzyme in the composition.

Background of the invention

Performance of a detergent product, for use in washing or cleaning method, such as a laundry or dishwashing method, is judged by a number of factors, including the ability to remove soils, and the ability to prevent the redeposition of the soils, or the breakdown products of the soils on the articles in the wash.

Food soils are often difficult to remove effectively from a soiled substrate. Highly coloured or 'dried-on' soils derived from fruit and/or vegetables are particularly challenging soils to remove. Specific examples of such soils would include orange, tomato, banana, tea, mango, broccoli soils and grass. Also residues from plants usual contaminating inorganic compounds like mud or clay are very difficult to remove. The substrates can be fabrics, or hard surfaces such as glassware or chinaware.

Xylanolytic enzymes like xylanase are commonly used in the paper/pulp industry and to a lesser extent also in the starch/baking industry. Examples of such xylanolytic enzymes are the commercial available xylanases e.g. Pulpzyme HB and SP431 (Novo Nordisk A/S), Lyxasan (Gist-Brocades) Optipulp and Xylanase (Solvay).

In WO94/01532 is disclosed a method for producing enzymes from strains of the alkalophilic species *Bacillus* sp. AC13. Enzymes obtainable from these strains are proteases, xylanases and cellulases as well. The protease and cellulase enzymes are demonstrated to be valuable for use in detergents while on the other hand xylanase is shown to be of use in processes for treatment of lignocellulosic pulp i.e. the paper pulp industry.

In WO92/06209 is disclosed that xylanase enzymes are overexpressed by microbial strains constructed via genetic techniques free of cellulolytic enzymes. These xylanase enzymes are described for use in a variety of applications like the bleaching of wood pulps and the modification of cereals and grains for use in baking and the production of animal feeds.

In WO92/19726 are disclosed stabilized, modified enzymes. The naturally occurring amino acids (other than proline) have been substituted with a proline residue at one or more positions. Among the modified enzymes, amylases,

lipases, cellulases, xylanases and peroxidases are mentioned. Said stabilized, modified enzymes can be used in detergent compositions.

However benefits for use of xylanolytic enzymes at low levels in cleaning formulations have not been previously recognized.

Summary of the invention

We have now found that xylanolytic enzymes may valuably be incorporated into cleaning compositions at a level of from 0.0001% to 2% of the composition by weight of active enzyme. The inclusion of such enzymes provides stain/soil removal benefits. Removal of food soils/stains, and in particular the removal of dried-on fruit, vegetable soils/stains and/or polysaccharides is enabled. We now have surprisingly found that application of xylanase at above-mentioned levels in cleaning compositions delivers an important boost of the cleaning performance.

It is an object of the present invention to provide cleaning compositions containing enzymes with xylanolytic activity and in particular to provide laundry and dishwashing detergent compositions containing said xylanolytic enzymes providing enhanced soil removal of fruit and/or vegetables stains.

Detailed description of the invention

An essential component of the cleaning compositions of the invention is a xylanolytic enzyme. This enzyme, preferably xylanase, is incorporated into the compositions in accordance with the invention at a level of from 0.0001% to 2%, preferably from 0.0005% to 0.5%, more preferably from 0.001% to 0.05% active enzyme by weight of the composition.

By xylanolytic enzyme it is meant herein any enzyme which degrade, for instance hydrolyze and/or modify, xylan containing polymers which are associated with hemicellulose and other plant polysaccharides.

Xylanolytic enzymes herein include naturally derived xylanolytic enzymes and any variants showing a positive immunological cross-reaction with an antibody directed against natural occurring xylanase.

Any such variants may be specifically designed with regard to the optimization of performance efficiency in the detergent compositions of the invention. For example, variants may be designed such that the compatibility of the enzyme to commonly encountered components of such compositions is increased.

Alternatively, the variant may be designed such that the optimal pH, bleach stability, catalytic activity and the like of the enzyme variant is tailored to suit the particular detergent application.

Xylanolytic enzymes include the commercial available xylanases like Pulpzyme HB and SP431 (Novo Nordisk), Lyxasan (Gist-Brocades), Optipulp and Xylanase (Solvay) as well.

Detergent components

The detergent compositions of the invention may also contain additional detergent components. The precise nature of these additional components, and levels of incorporation thereof will depend on the physical form of the composition, and the nature of the cleaning operation for which it is to be used.

The compositions of the invention may for example, be formulated as manual and machine dishwashing compositions, hand and machine laundry detergent compositions including laundry additive compositions and compositions suitable for

use in the pretreatment of stained fabrics, rinse added fabric softener compositions, and compositions for use in general household hard surface cleaning operations.

When formulated as compositions suitable for use in a machine washing method, the compositions of the invention preferably contain both a surfactant and a builder compound and additionally one or more detergent components preferably selected from organic polymeric compounds, bleaching agents, additional enzymes, suds suppressors, dispersants, lime-soap dispersants, soil suspension and anti-redeposition agents and corrosion inhibitors. Laundry compositions can also contain softening agents, as additional detergent components.

When formulated as compositions for use in manual dishwashing methods the compositions of the invention preferably contain a surfactant and preferably other detergent components selected from organic polymeric compounds, suds enhancing agents, group II metal ions, solvents, hydrotropes and additional enzymes.

If needed the density of the laundry detergent compositions herein ranges from 550 to 1000 g/liter, preferably 600 to 950 g/liter of composition measured at 20°C.

The "compact" form of the compositions herein is best reflected by density and, in terms of composition, by the amount of inorganic filler salt; inorganic filler salts are conventional ingredients of detergent compositions in powder form; in conventional detergent compositions, the filler salts are present in substantial amounts, typically 17-35% by weight of the total composition.

In the compact compositions, the filler salt is present in amounts not exceeding 15% of the total composition, preferably not exceeding 10%, most preferably not exceeding 5% by weight of the composition.

The inorganic filler salts, such as meant in the present compositions are selected from the alkali and alkaline-earth-metal salts of sulphates and chlorides.

A preferred filler salt is sodium sulphate.

Surfactant system

The detergent compositions according to the present invention comprise a surfactant system wherein the surfactant can be selected from nonionic and/or anionic and/or cationic and/or ampholytic and/or zwitterionic and/or semi-polar surfactants.

The surfactant is typically present at a level of from 0.1% to 60% by weight. More preferred levels of incorporation are 1% to 35% by weight, most preferably from 1% to 20% by weight of machine dishwashing, laundry and rinse added fabric softener compositions in accord with the invention, and from 5% to 60% by weight, more preferably from 15% to 45% by weight of manual dishwashing compositions in accord with the invention.

The surfactant is preferably formulated to be compatible with enzyme components present in the composition. In liquid or gel compositions the surfactant is most preferably formulated such that it promotes, or at least does not degrade, the stability of any enzyme in these compositions.

Preferred non-alkylbenzene sulfonate surfactant systems to be used according to the present invention comprise as a surfactant one or more of the nonionic and/or anionic surfactants described herein.

Polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols are suitable for use as the nonionic surfactant of the surfactant systems of the present invention, with the polyethylene oxide condensates being


preferred. These compounds include the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 14 carbon atoms, preferably from about 8 to about 14 carbon atoms, in either a straight-chain or branched-chain configuration with the alkylene oxide. In a preferred embodiment, the ethylene oxide is present in an amount equal to from about 2 to about 25 moles, more preferably from about 3 to about 15 moles, of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include IgepalTM CO-630, marketed by the GAF Corporation; and TritonTM X-45, X-114, X-100 and X-102, all marketed by the Rohm & Haas Company. These surfactants are commonly referred to as alkylphenol alkoxyates (e.g., alkyl phenol ethoxyates).

The condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethylene oxide are suitable for use as the nonionic surfactant of the nonionic surfactant systems of the present invention. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Preferred are the condensation products of alcohols having an alkyl group containing from about 8 to about 20 carbon atoms, more preferably from about 10 to about 18 carbon atoms, with from about 2 to about 10 moles of ethylene oxide per mole of alcohol. About 2 to about 7 moles of ethylene oxide and most preferably from 2 to 5 moles of ethylene oxide per mole of alcohol are present in said condensation products. Examples of commercially available nonionic surfactants of this type include TergitolTM 15-S-9 (the condensation product of C₁₁-C₁₅ linear alcohol with 9 moles ethylene oxide), TergitolTM 24-L-6 NMW (the condensation product of C₁₂-C₁₄ primary alcohol with 6 moles ethylene oxide with a narrow molecular weight distribution), both marketed by Union Carbide Corporation; NeodolTM 45-9 (the condensation product of C₁₄-C₁₅ linear alcohol with 9 moles of ethylene oxide), NeodolTM 23-3 (the condensation product of C₁₂-C₁₃ linear alcohol with

3.0 moles of ethylene oxide), NeodolTM 45-7 (the condensation product of C₁₄-C₁₅ linear alcohol with 7 moles of ethylene oxide), NeodolTM 45-5 (the condensation product of C₁₄-C₁₅ linear alcohol with 5 moles of ethylene oxide) marketed by Shell Chemical Company, KyroTM EOB (the condensation product of C₁₃-C₁₅ alcohol with 9 moles ethylene oxide), marketed by The Procter & Gamble Company, and Genapol LA 050 (the condensation product of C₁₂-C₁₄ alcohol with 5 moles of ethylene oxide) marketed by Hoechst. Preferred range of HLB in these products is from 8-11 and most preferred from 8-10.

Also useful as the nonionic surfactant of the surfactant systems of the present invention are the alkylpolysaccharides disclosed in U.S. Patent 4,565,647, Llenado, issued January 21, 1986, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g. a polyglycoside, hydrophilic group containing from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties (optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside). The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6- positions on the preceding saccharide units.

The preferred alkylpolyglycosides have the formula



wherein R² is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 2 or 3, preferably 2; t is from 0 to about 10,

preferably 0; and x is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4- and/or 6-position, preferably predominately the 2-position.


The condensation products of ethylene oxide with a hydrophobic base formed by the condensation of propylene oxide with propylene glycol are also suitable for use as the additional nonionic surfactant systems of the present invention. The hydrophobic portion of these compounds will preferably have a molecular weight of from about 1500 to about 1800 and will exhibit water insolubility. The addition of polyoxyethylene moieties to this hydrophobic portion tends to increase the water solubility of the molecule as a whole, and the liquid character of the product is retained up to the point where the polyoxyethylene content is about 50% of the total weight of the condensation product, which corresponds to condensation with up to about 40 moles of ethylene oxide. Examples of compounds of this type include certain of the commercially-available PluronicTM surfactants, marketed by BASF.

Also suitable for use as the nonionic surfactant of the nonionic surfactant system of the present invention, are the condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylenediamine. The hydrophobic moiety of these products consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from about 2500 to about 3000. This hydrophobic moiety is condensed with ethylene oxide to the extent that the condensation product contains from about 40% to about 80% by

weight of polyoxyethylene and has a molecular weight of from about 5,000 to about 11,000. Examples of this type of nonionic surfactant include certain of the commercially available TetronicTM compounds, marketed by BASF.

Preferred for use as the nonionic surfactant of the surfactant systems of the present invention are polyethylene oxide condensates of alkyl phenols, condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethylene oxide, alkylpolysaccharides, and mixtures thereof. Most preferred are C₈-C₁₄ alkyl phenol ethoxylates having from 3 to 15 ethoxy groups and C₈-C₁₈ alcohol ethoxylates (preferably C₁₀ avg.) having from 2 to 10 ethoxy groups, and mixtures thereof.

Highly preferred nonionic surfactants are polyhydroxy fatty acid amide surfactants of the formula.



wherein R¹ is H, or R¹ is C₁₋₄ hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl or a mixture thereof, R² is C₅₋₃₁ hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative thereof. Preferably, R¹ is methyl, R² is a straight C₁₁₋₁₅ alkyl or C₁₆₋₁₈ alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z is derived from a reducing sugar such as glucose, fructose, maltose, lactose, in a reductive amination reaction.


When included in such laundry detergent compositions, the nonionic surfactant systems of the present invention act to improve the greasy/oily stain removal properties of such laundry detergent compositions across a broad range of laundry conditions.

Highly preferred anionic surfactants include alkyl alkoxyated sulfate surfactants hereof are water soluble salts or acids of the formula $RO(A)_mSO_3M$ wherein R is an unsubstituted C_{10} - C_{24} alkyl or hydroxyalkyl group having a C_{10} - C_{24} alkyl component, preferably a C_{12} - C_{20} alkyl or hydroxyalkyl, more preferably C_{12} - C_{18} alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxyated sulfates as well as alkyl propoxyated sulfates are contemplated herein. Specific examples of substituted ammonium cations include methyl-, dimethyl, trimethyl-ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like. Exemplary surfactants are C_{12} - C_{18} alkyl polyethoxylate (1.0) sulfate (C_{12} - $C_{18}E(1.0)M$), C_{12} - C_{18} alkyl polyethoxylate (2.25) sulfate (C_{12} - $C_{18}E(2.25)M$), C_{12} - C_{18} alkyl polyethoxylate (3.0) sulfate (C_{12} - $C_{18}E(3.0)M$), and C_{12} - C_{18} alkyl polyethoxylate (4.0) sulfate (C_{12} - $C_{18}E(4.0)M$), wherein M is conveniently selected from sodium and potassium.

Suitable anionic surfactants to be used are alkyl ester sulfonate surfactants including linear esters of C_8 - C_{20} carboxylic acids (i.e., fatty acids) which are sulfonated with gaseous SO_3 according to "The Journal of the American Oil Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm oil, etc.

The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester sulfonate surfactants of the structural formula :

12



wherein R^3 is a C_8 - C_{20} hydrocarbyl, preferably an alkyl, or combination thereof, R^4 is a C_1 - C_6 hydrocarbyl, preferably an alkyl, or combination thereof, and M is a cation which forms a water soluble salt with the alkyl ester sulfonate. Suitable salt-forming cations include metals such as sodium, potassium, and lithium, and substituted or unsubstituted ammonium cations, such as monoethanolamine, diethanolamine, and triethanolamine. Preferably, R^3 is C_{10} - C_{16} alkyl, and R^4 is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates wherein R^3 is C_{10} - C_{16} alkyl.

Other suitable anionic surfactants include the alkyl sulfate surfactants which are water soluble salts or acids of the formula ROSO_3M wherein R preferably is a C_{10} - C_{24} hydrocarbyl, preferably an alkyl or hydroxyalkyl having a C_{10} - C_{20} alkyl component, more preferably a C_{12} - C_{18} alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium, potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like). Typically, alkyl chains of C_{12} - C_{16} are preferred for lower wash temperatures (e.g. below about 50°C) and C_{16} - 18 alkyl chains are preferred for higher wash temperatures (e.g. above about 50°C).

Other anionic surfactants useful for deterative purposes can also be included in the laundry detergent compositions of the present invention. These can include salts (including, for example, sodium, potassium, ammonium, and substituted

ammonium salts such as mono-, di- and triethanolamine salts) of soap, C₈-C₂₂ primary or secondary alkanesulfonates, C₈-C₂₄ olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, C₈-C₂₄ alkylpolyglycoethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated C₁₂-C₁₈ monoesters) and diesters of sulfosuccinates (especially saturated and unsaturated C₆-C₁₂ diesters), acyl sarcosinates, sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), branched primary alkyl sulfates, and alkyl polyethoxy carboxylates such as those of the formula $RO(CH_2CH_2O)_k-CH_2COO-M^+$ wherein R is a C₈-C₂₂ alkyl, k is an integer from 1 to 10, and M is a soluble salt-forming cation. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tall oil.


Further examples are described in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in U.S. Patent 3,929,678, issued December 30, 1975 to Laughlin, et al. at Column 23, line 58 through Column 29, line 23 (herein incorporated by reference).

When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to about 40%, preferably from about 3% to about 20% by weight of such anionic surfactants.

The laundry detergent compositions of the present invention may also contain cationic, ampholytic, zwitterionic, and


semi-polar surfactants, as well as the nonionic and/or anionic surfactants other than those already described herein.

Cationic deterative surfactants suitable for use in the laundry detergent compositions of the present invention are those having one long-chain hydrocarbyl group. Examples of such cationic surfactants include the ammonium surfactants such as alkyltrimethylammonium halogenides, and those surfactants having the formula :



wherein R^2 is an alkyl or alkyl benzyl group having from about 8 to about 18 carbon atoms in the alkyl chain, each R^3 is selected from the group consisting of $-CH_2CH_2-$, $-CH_2CH(CH_3)-$, $-CH_2CH(CH_2OH)-$, $-CH_2CH_2CH_2-$, and mixtures thereof; each R^4 is selected from the group consisting of C_1 - C_4 alkyl, C_1 - C_4 hydroxyalkyl, benzyl ring structures formed by joining the two R^4 groups, $-CH_2CHOH-CHOHCOR^6CHOHCH_2OH$ wherein R^6 is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when y is not 0; R^5 is the same as R^4 or is an alkyl chain wherein the total number of carbon atoms of R^2 plus R^5 is not more than about 18; each y is from 0 to about 10 and the sum of the y values is from 0 to about 15; and X is any compatible anion.

Highly preferred cationic surfactants are the water-soluble quaternary ammonium compounds useful in the present composition having the formula :



wherein R_1 is C_8 - C_{16} alkyl, each of R_2 , R_3 and R_4 is independently C_1 - C_4 alkyl, C_1 - C_4 hydroxy alkyl, benzyl, and $-(C_2H_4O)_xH$ where x has a value from 2 to 5, and X is an anion. Not more than one of R_2 , R_3 or R_4 should be benzyl.

The preferred alkyl chain length for R_1 is C_{12} - C_{15} particularly where the alkyl group is a mixture of chain lengths derived from coconut or palm kernel fat or is derived synthetically by olefin build up or OXO alcohols synthesis.

Preferred groups for R_2R_3 and R_4 are methyl and hydroxyethyl groups and the anion X may be selected from halide, methosulphate, acetate and phosphate ions.

Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are :

coconut trimethyl ammonium chloride or bromide;
 coconut methyl dihydroxyethyl ammonium chloride or bromide;
 decyl triethyl ammonium chloride;
 decyl dimethyl hydroxyethyl ammonium chloride or bromide;
 C_{12-15} dimethyl hydroxyethyl ammonium chloride or bromide;
 coconut dimethyl hydroxyethyl ammonium chloride or bromide;
 myristyl trimethyl ammonium methyl sulphate;
 lauryl dimethyl benzyl ammonium chloride or bromide;
 lauryl dimethyl (ethenoxy)₄ ammonium chloride or bromide;
 choline esters (compounds of formula (i) wherein R_1 is $CH_2-CH_2-O-C-C_{12-14}$ alkyl and $R_2R_3R_4$ are methyl).

||

O

di-alkyl imidazolines [compounds of formula (i)].

Other cationic surfactants useful herein are also described in U.S. Patent 4,228,044, Cambre, issued October 14, 1980 and in European Patent Application EP 000,224.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 25%, preferably from about 1% to about 8% by weight of such cationic surfactants.

Ampholytic surfactants are also suitable for use in the laundry detergent compositions of the present invention. These surfactants can be broadly described as aliphatic derivatives of secondary or tertiary amines, or aliphatic derivatives of heterocyclic secondary and tertiary amines in which the aliphatic radical can be straight- or branched-chain. One of the aliphatic substituents contains at least about 8 carbon atoms, typically from about 8 to about 18 carbon atoms, and at least one contains an anionic water-

solubilizing group, e.g. carboxy, sulfonate, sulfate. See U.S. Patent No. 3,929,678 to Laughlin et al., issued December 30, 1975 at column 19, lines 18-35, for examples of ampholytic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such ampholytic surfactants.


Zwitterionic surfactants are also suitable for use in laundry detergent compositions. These surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. See U.S. Patent No. 3,929,678 to Laughlin et al., issued December 30, 1975 at column 19, line 38 through column 22, line 48, for examples of zwitterionic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such zwitterionic surfactants.

Semi-polar nonionic surfactants are a special category of nonionic surfactants which include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; water-soluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and a moiety selected from the group

consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms.

Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula



wherein R^3 is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms; R^4 is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof; x is from 0 to about 3; and each R^5 is an alkyl or hydroxyalkyl group containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The R^5 groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.

These amine oxide surfactants in particular include C_{10} - C_{18} alkyl dimethyl amine oxides and C_8 - C_{12} alkoxy ethyl dihydroxy ethyl amine oxides.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such semi-polar nonionic surfactants.

Optional detergent ingredients :

Preferred detergent compositions of the present invention may further comprise an enzyme which provides cleaning performance and/or fabric care benefits. Said enzymes include enzymes selected from cellulases, hemicellulases, peroxidases, proteases, gluco-amylases, amylases, lipases, cutinases, pectinases, reductases, oxidases, phenoloxidases, lipxygenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases or mixtures thereof.

A preferred combination is a cleaning composition having cocktail of conventional applicable enzymes like protease, amylase, lipase, cutinase and/or cellulase in conjunction with one or more plant cell wall degrading enzymes.

The cellulases usable in the present invention include both bacterial or fungal cellulase. Preferably, they will have a pH optimum of between 5 and 9.5. Suitable cellulases are disclosed in U.S. Patent 4,435,307, Barbesgaard et al, which discloses fungal cellulase produced from *Humicola insolens*. Suitable cellulases are also disclosed in GB-A-2.075.028; GB-A-2.095.275 and DE-OS-2.247.832.

Examples of such cellulases are cellulases produced by a strain of *Humicola insolens* (*Humicola grisea* var. *thermoidea*), particularly the *Humicola* strain DSM 1800. Other suitable cellulases are cellulases originated from *Humicola insolens* having a molecular weight of about 50KDa, an isoelectric point of 5.5 and containing 415 amino acids. Especially suitable cellulases are the cellulases having color care benefits. Examples of such cellulases are cellulases described in European patent application No. 91202879.2, filed November 6, 1991 (Novo).

Peroxidase enzymes are used in combination with oxygen sources, e.g. percarbonate, perborate, persulfate, hydrogen peroxide, etc. They are used for "solution bleaching", i.e. to prevent transfer of dyes or pigments removed from substrates during wash operations to other substrates in the wash solution. Peroxidase enzymes are known in the art, and include, for example, horseradish peroxidase, ligninase, and haloperoxidase such as chloro- and bromo-peroxidase. Peroxidase-containing detergent compositions are disclosed, for example, in PCT International Application WO 89/099813 and in European Patent application EP No. 91202882.6, filed on November 6, 1991.

Said cellulases and/or peroxidases are normally incorporated in the detergent composition at levels from

0.0001% to 2% of active enzyme by weight of the detergent composition.

Preferred commercially available protease enzymes include those sold under the tradenames Alcalase, Savinase, Primase, Durazym, and Esperase by Novo Nordisk A/S (Denmark), those sold under the tradename Maxatase, Maxacal and Maxapem by Gist-Brocades, those sold by Genencor International, and those sold under the tradename Opticlean and Optimase by Solvay Enzymes. Also proteases described in our co-pending application USSN 08/136,797 can be included in the detergent composition of the invention. Protease enzyme may be incorporated into the compositions in accordance with the invention at a level of from 0.0001% to 2% active enzyme by weight of the composition.

Other preferred enzymes that can be included in the detergent compositions of the present invention include lipases. Suitable lipase enzymes for detergent usage include those produced by microorganisms of the *Pseudomonas* group, such as *Pseudomonas stutzeri* ATCC 19.154, as disclosed in British Patent 1,372,034. Suitable lipases include those which show a positive immunological cross-reaction with the antibody of the lipase, produced by the microorganism *Pseudomonas fluorescent* IAM 1057. This lipase is available from Amano Pharmaceutical Co. Ltd., Nagoya, Japan, under the trade name Lipase P "Amano," hereinafter referred to as "Amano-P". Especially suitable lipases are lipases such as M1 Lipase^R and Lipomax^R (Gist-Brocades) and Lipolase^R (Novo) which have found to be very effective when used in combination with the compositions of the present invention. Also suitable are cutinases [EC 3.1.1.50] which can be considered as a special kind of lipase, namely lipases which do not require interfacial activation. Addition of cutinases to detergent compositions have been described in e.g. WO-A-88/09367 (Genencor).

The lipases and/or cutinases are normally incorporated in the detergent composition at levels from 0.0001% to 2% of active enzyme by weight of the detergent composition.

Amylases (α and/or β) can be included for removal of carbohydrate-based stains. Suitable amylases are Termamyl^R (Novo Nordisk), Fungamyl^R and BAN^R (Novo Nordisk).

The above-mentioned enzymes may be of any suitable origin, such as vegetable, animal, bacterial, fungal and yeast origin.

Said enzymes are normally incorporated in the detergent composition at levels from 0.0001% to 2% of active enzyme by weight of the detergent composition.

Other suitable detergent ingredients that can be added are enzyme oxidation scavengers which are described in Copending European Patent application 92870018.6 filed on January 31, 1992. Examples of such enzyme oxidation scavengers are ethoxylated tetraethylene polyamines.

Especially preferred detergent ingredients are combinations with technologies which also provide a type of color care benefit. Examples of these technologies are metallo catalysts for color maintenance. Such metallo catalysts are described in copending European Patent Application No. 92870181.2.

Additional optional detergent ingredients that can be included in the detergent compositions of the present invention include bleaching agents such as PB1, PB4 and percarbonate with a particle size of 400-800 microns. These bleaching agent components can include one or more oxygen bleaching agents and, depending upon the bleaching agent chosen, one or more bleach activators. When present oxygen bleaching compounds will typically be present at levels of from about 1% to about 25%. In general, bleaching compounds are optional components in non-liquid formulations, e.g. granular detergents.

The bleaching agent component for use herein can be any of the bleaching agents useful for detergent compositions including oxygen bleaches as well as others known in the art. The bleaching agent suitable for the present invention can be an activated or non-activated bleaching agent.

One category of oxygen bleaching agent that can be used encompasses percarboxylic acid bleaching agents and salts thereof. Suitable examples of this class of agents include magnesium monoperoxyphthalate hexahydrate, the magnesium salt of meta-chloro perbenzoic acid, 4-nonylamino-4-oxoperoxybutyric acid and diperoxydodecanedioic acid. Such bleaching agents are disclosed in U.S. Patent 4,483,781, U.S. Patent Application 740,446, European Patent Application 0,133,354 and U.S. Patent 4,412,934. Highly preferred bleaching agents also include 6-nonylamino-6-oxoperoxypropionic acid as described in U.S. Patent 4,634,551.

Another category of bleaching agents that can be used encompasses the halogen bleaching agents. Examples of hypohalite bleaching agents, for example, include trichloro isocyanuric acid and the sodium and potassium dichloroisocyanurates and N-chloro and N-bromo alkane sulphonamides. Such materials are normally added at 0.5-10% by weight of the finished product, preferably 1-5% by weight.

The hydrogen peroxide releasing agents can be used in combination with bleach activators such as tetraacetylenediamine (TAED), nonanoyloxybenzenesulfonate (NOBS, described in US 4,412,934), 3,5-trimethylhexanoyloxybenzenesulfonate (ISONOBS, described in EP 120,591) or pentaacetylglucose (PAG), which are perhydrolyzed to form a peracid as the active bleaching species, leading to improved bleaching effect. Also suitable activators are acylated citrate esters such as disclosed in Copending European Patent Application No. 91870207.7.

Useful bleaching agents, including peroxyacids and bleaching systems comprising bleach activators and peroxygen

bleaching compounds for use in cleaning compositions according to the invention are described in our co-pending application USSN 08/136,626.

The hydrogen peroxide may also be present by adding an enzymatic system (i.e. an enzyme and a substrate therefore) which is capable of generating hydrogen peroxide at the beginning or during the washing and/or rinsing process. Such enzymatic systems are disclosed in EP Patent Application 91202655.6 filed October 9, 1991.

Bleaching agents other than oxygen bleaching agents are also known in the art and can be utilized herein. One type of non-oxygen bleaching agent of particular interest includes photoactivated bleaching agents such as the sulfonated zinc and/or aluminum phthalocyanines. These materials can be deposited upon the substrate during the washing process. Upon irradiation with light, in the presence of oxygen, such as by hanging clothes out to dry in the daylight, the sulfonated zinc phthalocyanine is activated and, consequently, the substrate is bleached. Preferred zinc phthalocyanine and a photoactivated bleaching process are described in U.S. Patent 4,033,718. Typically, detergent compositions will contain about 0.025% to about 1.25%, by weight, of sulfonated zinc phthalocyanine.

The compositions according to the present invention may further comprise a builder system. Any conventional builder system is suitable for use herein including aluminosilicate materials, silicates, polycarboxylates and fatty acids, materials such as ethylenediamine tetraacetate, metal ion sequestrants such as aminopolyphosphonates, particularly ethylenediamine tetramethylene phosphonic acid and diethylene triamine pentamethylenephosphonic acid. Though less preferred for obvious environmental reasons, phosphate builders can also be used herein.

Suitable builders can be an inorganic ion exchange material, commonly an inorganic hydrated aluminosilicate material, more particularly a hydrated synthetic zeolite such as hydrated zeolite A, X, B, HS or MAP.

Another suitable inorganic builder material is layered silicate, e.g. SKS-6 (Hoechst). SKS-6 is a crystalline layered silicate consisting of sodium silicate ($\text{Na}_2\text{Si}_2\text{O}_5$).

Suitable polycarboxylates containing one carboxy group include lactic acid, glycolic acid and ether derivatives thereof as disclosed in Belgian Patent Nos. 831,368, 821,369 and 821,370. Polycarboxylates containing two carboxy groups include the water-soluble salts of succinic acid, malonic acid, (ethylenedioxy) diacetic acid, maleic acid, diglycollic acid, tartaric acid, tartronic acid and fumaric acid, as well as the ether carboxylates described in German Offenlegenschrift 2,446,686, and 2,446,687 and U.S. Patent No. 3,935,257 and the sulfinyl carboxylates described in Belgian Patent No. 840,623. Polycarboxylates containing three carboxy groups include, in particular, water-soluble citrates, aconitrates and citraconates as well as succinate derivatives such as the carboxymethyloxysuccinates described in British Patent No. 1,379,241, lactoxysuccinates described in Netherlands Application 7205873, and the oxypolycarboxylate materials such as 2-oxa-1,1,3-propane tricarboxylates described in British Patent No. 1,387,447.

Polycarboxylates containing four carboxy groups include oxydisuccinates disclosed in British Patent No. 1,261,829, 1,1,2,2-ethane tetracarboxylates, 1,1,3,3-propane tetracarboxylates and 1,1,2,3-propane tetracarboxylates. Polycarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in British Patent Nos. 1,398,421 and 1,398,422 and in U.S. Patent No. 3,936,448, and the sulfonated pyrolysed citrates described in British Patent No. 1,082,179, while polycarboxylates containing phosphone substituents are disclosed in British Patent No. 1,439,000.

Alicyclic and heterocyclic polycarboxylates include cyclopentane-cis,cis,cis-tetracarboxylates, cyclopentadienide pentacarboxylates, 2,3,4,5-tetrahydro-furan - cis, cis, cis-tetracarboxylates, 2,5-tetrahydro-furan -cis -dicarboxylates, 2,2,5,5-tetrahydrofuran - tetracarboxylates, 1,2,3,4,5,6-hexane -hexacarboxylates and and carboxymethyl derivatives of polyhydric alcohols such as sorbitol, mannitol and xylitol. Aromatic poly-carboxylates include mellitic acid, pyromellitic acid and the phthalic acid derivatives disclosed in British Patent No. 1,425,343.

Of the above, the preferred polycarboxylates are hydroxycarboxylates containing up to three carboxy groups per molecule, more particularly citrates.

Preferred builder systems for use in the present compositions include a mixture of a water-insoluble aluminosilicate builder such as zeolite A or of a layered silicate (SKS-6), and a water-soluble carboxylate chelating agent such as citric acid.

A suitable chelant for inclusion in the detergent compositions in accordance with the invention is ethylenediamine-N,N'-disuccinic acid (EDDS) or the alkali metal, alkaline earth metal, ammonium, or substituted ammonium salts thereof, or mixtures thereof. Preferred EDDS compounds are the free acid form and the sodium or magnesium salt thereof. Examples of such preferred sodium salts of EDDS include Na_2EDDS and Na_4EDDS . Examples of such preferred magnesium salts of EDDS include MgEDDS and Mg_2EDDS . The magnesium salts are the most preferred for inclusion in compositions in accordance with the invention.

Preferred builder systems include a mixture of a water-insoluble aluminosilicate builder such as zeolite A, and a watersoluble carboxylate chelating agent such as citric acid. Other builder materials that can form part of the builder system for use in granular compositions include inorganic materials such as alkali metal carbonates, bicarbonates,

silicates, and organic materials such as the organic phosphonates, amino polyalkylene phosphonates and amino polycarboxylates.

Other suitable water-soluble organic salts are the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms. Polymers of this type are disclosed in GB-A-1,596,756. Examples of such salts are polyacrylates of MW 2000-5000 and their copolymers with maleic anhydride, such copolymers having a molecular weight of from 20,000 to 70,000, especially about 40,000.

Detergency builder salts are normally included in amounts of from 10% to 80% by weight of the composition preferably from 20% to 70% and most usually from 30% to 60% by weight.

Another optional ingredient is a suds suppressor, exemplified by silicones, and silica-silicone mixtures. Silicones can be generally represented by alkylated polysiloxane materials while silica is normally used in finely divided forms exemplified by silica aerogels and xerogels and hydrophobic silicas of various types. These materials can be incorporated as particulates in which the suds suppressor is advantageously releasably incorporated in a water-soluble or water-dispersible, substantially non-surface-active detergent impermeable carrier. Alternatively the suds suppressor can be dissolved or dispersed in a liquid carrier and applied by spraying on to one or more of the other components.

A preferred silicone suds controlling agent is disclosed in Bartollota et al. U.S. Patent 3 933 672. Other particularly useful suds suppressors are the self-emulsifying silicone suds suppressors, described in German Patent Application DTOS 2 646 126 published April 28, 1977. An example of such a compound is DC-544, commercially available

from Dow Corning, which is a siloxane-glycol copolymer. Especially preferred suds controlling agent are the suds suppressor system comprising a mixture of silicone oils and 2-alkyl-alkanols. Suitable 2-alkyl-alkanols are 2-butyl-octanol which are commercially available under the trade name Isofol 12 R.

Such suds suppressor system are described in Copending European Patent application N 92870174.7 filed 10 November, 1992.

Especially preferred silicone suds controlling agents are described in Copending European Patent application N°92201649.8. Said compositions can comprise a silicone/silica mixture in combination with fumed nonporous silica such as Aerosil^R.

The suds suppressors described above are normally employed at levels of from 0.001% to 2% by weight of the composition, preferably from 0.01% to 1% by weight.

Other components used in detergent compositions may be employed, such as soil-suspending agents, soil-release agents, optical brighteners, abrasives, bactericides, tarnish inhibitors, coloring agents, and/or encapsulated or non-encapsulated perfumes.

Especially suitable encapsulating materials are water soluble capsules which consist of a matrix of polysaccharide and polyhydroxy compounds such as described in GB 1,464,616.

Other suitable water soluble encapsulating materials comprise dextrans derived from ungelatinized starch acid-esters of substituted dicarboxylic acids such as described in US 3,455,838. These acid-ester dextrans are, preferably, prepared from such starches as waxy maize, waxy sorghum, sago, tapioca and potato. Suitable examples of said encapsulating materials include N-Lok manufactured by National Starch. The N-Lok encapsulating material consists of a modified maize starch and glucose. The starch is modified

by adding monofunctional substituted groups such as octenyl succinic acid anhydride.


Antiredeposition and soil suspension agents suitable herein include cellulose derivatives such as methylcellulose, carboxymethylcellulose and hydroxyethylcellulose, and homo- or co-polymeric polycarboxylic acids or their salts. Polymers of this type include the polyacrylates and maleic anhydride-acrylic acid copolymers previously mentioned as builders, as well as copolymers of maleic anhydride with ethylene, methylvinyl ether or methacrylic acid, the maleic anhydride constituting at least 20 mole percent of the copolymer. These materials are normally used at levels of from 0.5% to 10% by weight, more preferably from 0.75% to 8%, most preferably from 1% to 6% by weight of the composition.

Preferred optical brighteners are anionic in character, examples of which are disodium 4,4'-bis-(2-diethanolamino-4-anilino -s- triazin-6-ylamino)stilbene-2:2' disulphonate, disodium 4, - 4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino-stilbene-2:2' - disulphonate, disodium 4,4' - bis-(2,4-dianilino-s-triazin-6-ylamino)stilbene-2:2' - disulphonate, monosodium 4',4'' -bis-(2,4-dianilino-s-triazin-6 ylamino)stilbene-2-sulphonate, disodium 4,4' -bis-(2-anilino-4-(N-methyl-N-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2' - disulphonate, di-sodium 4,4' -bis-(4-phenyl-2,1,3-triazol-2-yl)-stilbene-2,2' disulphonate, di-sodium 4,4'bis(2-anilino-4-(1-methyl-2-hydroxyethylamino)-s-triazin-6-ylami-no)stilbene-2,2'disulphonate, sodium 2(stilbyl-4''-(naphtho-1',2':4,5)-1,2,3 - triazole-2''-sulphonate and 4,4'-bis(2-sulphostyryl)biphenyl.

Other useful polymeric materials are the polyethylene glycols, particularly those of molecular weight 1000-10000, more particularly 2000 to 8000 and most preferably about 4000. These are used at levels of from 0.20% to 5% more preferably from 0.25% to 2.5% by weight. These polymers and the previously mentioned homo- or co-polymeric

polycarboxylate salts are valuable for improving whiteness maintenance, fabric ash deposition, and cleaning performance on clay, proteinaceous and oxidizable soils in the presence of transition metal impurities.

Soil release agents useful in compositions of the present invention are conventionally copolymers or terpolymers of terephthalic acid with ethylene glycol and/or propylene glycol units in various arrangements. Examples of such polymers are disclosed in the commonly assigned US Patent Nos. 4116885 and 4711730 and European Published Patent Application No. 0 272 033. A particular preferred polymer in accordance with EP-A-0 272 033 has the formula



where PEG is $-(\text{OC}_2\text{H}_4)_n\text{O}-$, PO is $(\text{OC}_3\text{H}_6\text{O})$ and T is $(\text{pCOC}_6\text{H}_4\text{CO})$.

Also very useful are modified polyesters as random copolymers of dimethyl terephthalate, dimethyl sulfoisophthalate, ethylene glycol and 1-2 propane diol, the end groups consisting primarily of sulphobenzoate and secondarily of mono esters of ethylene glycol and/or propane diol. The target is to obtain a polymer capped at both end by sulphobenzoate groups, "primarily", in the present context most of said copolymers herein will be end-capped by sulphobenzoate groups. However, some copolymers will be less than fully capped, and therefore their end groups may consist of monoester of ethylene glycol and/or propane 1-2 diol, thereof consist "secondarily" of such species.

The selected polyesters herein contain about 46% by weight of dimethyl terephthalic acid, about 16% by weight of propane -1.2 diol, about 10% by weight ethylene glycol about 13% by weight of dimethyl sulfo benzoic acid and about 15% by weight of sulfoisophthalic acid, and have a molecular weight

of about 3.000. The polyesters and their method of preparation are described in detail in EPA 311 342.

Softening agents

Fabric softening agents can also be incorporated into laundry detergent compositions in accordance with the present invention. These agents may be inorganic or organic in type. Inorganic softening agents are exemplified by the smectite clays disclosed in GB-A-1 400 898 and in USP 5,019,292. Organic fabric softening agents include the water insoluble tertiary amines as disclosed in GB-A1 514 276 and EP-B0 011 340 and their combination with mono C12-C14 quaternary ammonium salts are disclosed in EP-B-0 026 527 and EP-B-0 026 528 and di-long-chain amides as disclosed in EP-B-0 242 919. Other useful organic ingredients of fabric softening systems include high molecular weight polyethylene oxide materials as disclosed in EP-A-0 299 575 and 0 313 146.

Levels of smectite clay are normally in the range from 5% to 15%, more preferably from 8% to 12% by weight, with the material being added as a dry mixed component to the remainder of the formulation. Organic fabric softening agents such as the water-insoluble tertiary amines or dilong chain amide materials are incorporated at levels of from 0.5% to 5% by weight, normally from 1% to 3% by weight whilst the high molecular weight polyethylene oxide materials and the water soluble cationic materials are added at levels of from 0.1% to 2%, normally from 0.15% to 1.5% by weight. These materials are normally added to the spray dried portion of the composition, although in some instances it may be more convenient to add them as a dry mixed particulate, or spray them as molten liquid on to other solid components of the composition.

Dye transfer inhibition

The present invention also relates to a process for inhibiting dye transfer from one fabric to another of solubilized and suspended dyes encountered during fabric laundering operations involving colored fabrics.


Polymeric dye transfer inhibiting agents

The detergent compositions according to the present invention also comprise from 0.001% to 10 %, preferably from 0.01% to 2%, more preferably from 0.05% to 1% by weight of polymeric dye transfer inhibiting agents. Said polymeric dye transfer inhibiting agents are normally incorporated into detergent compositions in order to inhibit the transfer of dyes from colored fabrics onto fabrics washed therewith. These polymers have the ability to complex or adsorb the fugitive dyes washed out of dyed fabrics before the dyes have the opportunity to become attached to other articles in the wash. Especially suitable polymeric dye transfer inhibiting agents are polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylpyrrolidone polymers, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof.


Addition of such polymers also enhances the performance of the enzymes according the invention.

a) Polyamine N-oxide polymers

The polyamine N-oxide polymers suitable for use contain units having the following structure formula :




wherein P is a polymerisable unit, whereto the R-N-O group can be attached to or wherein the R-N-O group forms part of the polymerisable unit or a combination of both.



A is NC, CO, C, -O-, -S-, -N- ; x is 0 or 1;

R are aliphatic, ethoxylated aliphatics, aromatic, heterocyclic or alicyclic groups or any combination thereof whereto the nitrogen of the N-O group can be attached or wherein the nitrogen of the N-O group is part of these groups.

The N-O group can be represented by the following general structures :



wherein R1, R2, and R3 are aliphatic groups, aromatic, heterocyclic or alicyclic groups or combinations thereof, x or/and y or/and z is 0 or 1 and wherein the nitrogen of the N-O group can be attached or wherein the nitrogen of the N-O group forms part of these groups.

The N-O group can be part of the polymerisable unit (P) or can be attached to the polymeric backbone or a combination of both.

Suitable polyamine N-oxides wherein the N-O group forms part of the polymerisable unit comprise polyamine N-oxides wherein R is selected from aliphatic, aromatic, alicyclic or heterocyclic groups.

One class of said polyamine N-oxides comprises the group of polyamine N-oxides wherein the nitrogen of the N-O group forms

part of the R-group. Preferred polyamine N-oxides are those wherein R is a heterocyclic group such as pyrridine, pyrrole, imidazole, pyrrolidine, piperidine, quinoline, acridine and derivatives thereof.

Another class of said polyamine N-oxides comprises the group of polyamine N-oxides wherein the nitrogen of the N-O group is attached to the R-group.

Other suitable polyamine N-oxides are the polyamine oxides whereto the N-O group is attached to the polymerisable unit.

Preferred class of these polyamine N-oxides are the polyamine N-oxides having the general formula (I) wherein R is an aromatic, heterocyclic or alicyclic groups wherein the nitrogen of the N-O functional group is part of said R group. Examples of these classes are polyamine oxides wherein R is a heterocyclic compound such as pyrridine, pyrrole, imidazole and derivatives thereof.

Another preferred class of polyamine N-oxides are the polyamine oxides having the general formula (I) wherein R are aromatic, heterocyclic or alicyclic groups wherein the nitrogen of the N-O functional group is attached to said R groups.

Examples of these classes are polyamine oxides wherein R groups can be aromatic such as phenyl.

Any polymer backbone can be used as long as the amine oxide polymer formed is water-soluble and has dye transfer inhibiting properties. Examples of suitable polymeric backbones are polyvinyls, polyalkylenes, polyesters, polyethers, polyamide, polyimides, polyacrylates and mixtures thereof.

The amine N-oxide polymers of the present invention typically have a ratio of amine to the amine N-oxide of 10:1 to 1:1000000. However the amount of amine oxide groups present in the polyamine oxide polymer can be varied by appropriate copolymerization or by appropriate degree of N-oxidation.

Preferably, the ratio of amine to amine N-oxide is from 2:3 to 1:1000000. More preferably from 1:4 to 1:1000000, most preferably from 1:7 to 1:1000000. The polymers of the present invention actually encompass random or block copolymers where one monomer type is an amine N-oxide and the other monomer type is either an amine N-oxide or not. The amine oxide unit of the polyamine N-oxides has a $\text{PKa} < 10$, preferably $\text{PKa} < 7$, more preferred $\text{PKa} < 6$.

The polyamine oxides can be obtained in almost any degree of polymerisation. The degree of polymerisation is not critical provided the material has the desired water-solubility and dye-suspending power.

Typically, the average molecular weight is within the range of 500 to 1000,000; preferably from 1,000 to 50,000, more preferably from 2,000 to 30,000, most preferably from 3,000 to 20,000.

b) Copolymers of N-vinylpyrrolidone and N-vinylimidazole

The N-vinylimidazole N-vinylpyrrolidone polymers used in the present invention have an average molecular weight range from 5,000-1,000,000, preferably from 20,000-200,000.

Highly preferred polymers for use in detergent compositions according to the present invention comprise a polymer selected from N-vinylimidazole N-vinylpyrrolidone copolymers wherein said polymer has an average molecular weight range from 5,000 to 50,000 more preferably from 8,000 to 30,000, most preferably from 10,000 to 20,000.

The average molecular weight range was determined by light scattering as described in Barth H.G. and Mays J.W. Chemical Analysis Vol 113, "Modern Methods of Polymer Characterization".

Highly preferred N-vinylimidazole N-vinylpyrrolidone copolymers have an average molecular weight range from 5,000 to 50,000; more preferably from 8,000 to 30,000; most preferably from 10,000 to 20,000.

The N-vinylimidazole N-vinylpyrrolidone copolymers characterized by having said average molecular weight range provide excellent dye transfer inhibiting properties while not adversely affecting the cleaning performance of detergent compositions formulated therewith.

The N-vinylimidazole N-vinylpyrrolidone copolymer of the present invention has a molar ratio of N-vinylimidazole to N-vinylpyrrolidone from 1 to 0.2, more preferably from 0.8 to 0.3, most preferably from 0.6 to 0.4 .

c) Polyvinylpyrrolidone

The detergent compositions of the present invention may also utilize polyvinylpyrrolidone ("PVP") having an average molecular weight of from about 2,500 to about 400,000, preferably from about 5,000 to about 200,000, more preferably from about 5,000 to about 50,000, and most preferably from about 5,000 to about 15,000. Suitable polyvinylpyrrolidones are commercially available from ISP Corporation, New York, NY and Montreal, Canada under the product names PVP K-15 (viscosity molecular weight of 10,000), PVP K-30 (average molecular weight of 40,000), PVP K-60 (average molecular weight of 160,000), and PVP K-90 (average molecular weight of 360,000). Other suitable polyvinylpyrrolidones which are commercially available from BASF Cooperation include Sokalan HP 165 and Sokalan HP 12; polyvinylpyrrolidones known to persons skilled in the detergent field (see for example EP-A-262,897 and EP-A-256,696).

d) Polyvinylloxazolidone :

The detergent compositions of the present invention may also utilize polyvinylloxazolidone as a polymeric dye transfer inhibiting agent. Said polyvinylloxazolidones have an average molecular weight of from about 2,500 to about 400,000, preferably from about 5,000 to about 200,000, more preferably from about 5,000 to about 50,000, and most preferably from about 5,000 to about 15,000.

e) Polyvinylimidazole :

The detergent compositions of the present invention may also utilize polyvinylimidazole as polymeric dye transfer inhibiting agent. Said polyvinylimidazoles have an average about 2,500 to about 400,000, preferably from about 5,000 to about 200,000, more preferably from about 5,000 to about 50,000, and most preferably from about 5,000 to about 15,000.

Method of washing

The process described herein comprises contacting fabrics with a laundering solution in the usual manner and exemplified hereunder.

The process of the invention is conveniently carried out in the course of the cleaning process. The method of cleaning is preferably carried out at 5 °C to 95 °C, especially between 10°C and 60°C. The pH of the treatment solution is preferably from 7 to 11, especially from 7.5 to 10.5.

The following examples are meant to exemplify compositions of the present invention, but are not necessarily meant to limit or otherwise define the scope of the invention.

In the detergent compositions, the abbreviated component identifications have the following meanings:

LAS	: Sodium linear C ₁₂ alkyl benzene sulphonate
TAS	: Sodium tallow alkyl sulphate
XYAS	: Sodium C _{1X} - C _{1Y} alkyl sulfate
SAS	: C ₁₂ -C ₁₄ secondary (2,3) alkyl sulfate in the form of the sodium salt.

- APG : Alkyl polyglycoside surfactant of formula $C_{12} - (\text{glycosyl})_x$, where x is 1.5,
- AEC : Alkyl ethoxycarboxylate surfactant of formula C_{12} ethoxy (2) carboxylate.
- SS : Secondary soap surfactant of formula 2-butyl octanoic acid
- 25EY : A C_{12} - C_{15} predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide
- 45EY : A C_{14} - C_{15} predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide
- XYEVS : C_{1X} - C_{1Y} sodium alkyl sulfate condensed with an average of Z moles of ethylene oxide per mole
- Nonionic : C_{13} - C_{15} mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5 sold under the tradename Plurafax LF404 by BASF GmbH
- CFAA : C_{12} - C_{14} alkyl N-methyl glucamide
- TFAA : C_{16} - C_{18} alkyl N-methyl glucamide.
- Silicate : Amorphous Sodium Silicate ($\text{SiO}_2:\text{Na}_2\text{O}$ ratio = 2.0)

NaSKS-6	: Crystalline layered silicate of formula Ω -Na ₂ Si ₂ O ₅
Carbonate	: Anhydrous sodium carbonate
Phosphate	: Sodium tripolyphosphate
MA/AA	: Copolymer of 1:4 maleic/acrylic acid, average molecular weight about 80,000
Polyacrylate	: Polyacrylate homopolymer with an average molecular weight of 8,000 sold under the tradename PA30 by BASF GmbH
Zeolite A	: Hydrated Sodium Aluminosilicate of formula Na ₁₂ (AlO ₂ SiO ₂) ₁₂ . 27H ₂ O having a primary particle size in the range from 1 to 10 micrometers
Citrate	: Tri-sodium citrate dihydrate
Citric	: Citric Acid
Perborate	: Anhydrous sodium perborate monohydrate bleach, empirical formula NaBO ₂ .H ₂ O ₂
PB4	: Anhydrous sodium perborate tetrahydrate
Percarbonate	: Anhydrous sodium percarbonate bleach of empirical formula 2Na ₂ CO ₃ .3H ₂ O ₂
TAED	: Tetraacetyl ethylene diamine

Paraffin	: Paraffin oil sold under the tradename Winog 70 by Wintershall.
Xylanase	: Xylanolytic enzyme sold under the tradenames Pulpzyme HB or SP431 by Novo Nordisk A/S or Lyxasan (Gist-Brocades) or Optipulp or Xylanase (Solvay).
Protease	: Proteolytic enzyme sold under the tradename Savinase by Novo Nordisk A/S.
Amylase	: Amylolytic enzyme sold under the tradename Termamyl by Novo Nordisk A/S
Lipase	: Lipolytic enzyme sold under the tradename Lipolase by Novo Nordisk A/S
Peroxidase	: Peroxidase enzyme
Cellulase	: Cellulosic enzyme sold under the tradename Carezyme or Celluzyme by Novo Nordisk A/S.
CMC	: Sodium carboxymethyl cellulose
HEDP	: 1,1-hydroxyethane diphosphonic acid
DETPMP	: Diethylene triamine penta (methylene phosphonic acid), marketed by Monsanto under the Trade name Dequest 2060
PVP	: Polyvinyl pyrrolidone polymer

EDDS	: Ethylenediamine -N, N'- disuccinic acid, [S,S] isomer in the form of the sodium salt.
Suds Suppressor	: 25% paraffin wax Mpt 50°C, 17% hydrophobic silica, 58% paraffin oil.
Granular Suds Suppressor	: 12% Silicone/silica, 18% stearyl alcohol, 70% starch in granular form
SCS	: Sodium cumene sulphonate
Sulphate	: Anhydrous sodium sulphate.
HMWPEO	: High molecular weight polyethylene oxide
PGMS	: Polyglycerol monostearate having a tradename of Radiesurf 248
TAE 25	: Tallow alcohol ethoxylate (25)

Assay for Xylanolytic Activity

A way to measure the xylanolytic activity is the response in endo-xylanase units (EXU), determined at pH 9.0 with remazol-xylan as substrate.

A xylanase sample is incubated with remazol-xylan substrate. The background of non-degraded dyed substrate is precipitated by ethanol. The remaining blue colour in the supernatant is proportional to the xylanase activity, and the xylanase units are then determined relatively to an enzyme standard at standard reaction conditions, i.e. at 50.0 +/- 0.1°C, pH 9.0, and 30 minutes reaction time.

In the following examples all levels of enzyme quoted are expressed as % active enzyme by weight of the composition:

Example 1

The following machine dishwashing detergent compositions were prepared (parts by weight) in accord with the invention.

	A	B	C	D	E	F
Citrate	24.0	-	-	24.0	24.0	29.0
Phosphate	-	30.0	46.0	-	-	-
MA/AA	6.0	-	-	6.0	6.0	-
Silicate	27.5	-	33.0	27.5	27.5	25.7
Carbonate	12.5	23.5	-	12.5	12.5	-
Perborate	10.4	10.4	10.4	10.4	10.4	1.9
PB4	-	-	-	-	-	8.7
TAED	3.0	3.0	3.0	3.0	3.0	4.4
Benzotriazole	-	0.3	-	-	-	0.3
Paraffin	-	0.5	-	-	-	0.5
HEDP	-	-	-	-	-	0.5
Protease	0.04	0.04	0.04	0.04	0.04	0.04
Amylase	0.02	0.01	0.01	0.02	0.01	0.02
Lipase	0.03	-	0.03	0.03	0.03	-
Xylanase	0.05	0.07	0.04	0.01	0.08	0.05
Pectinase	0.02	-	0.08	-	0.01	0.04
Nonionic	-	1.5	1.5	1.5	1.5	1.5
Sulphate	1.4	2.4	2.4	12.1	12.1	3.0
35AE3S	-	-	5.0	-	5.0	-
Granular Suds	1.0	-	-	-	-	-
Suppressor						
Water & minors				Up to 100%		

The compositions provide good soil removal when used in a machine dishwashing process.

Example 2

The following liquid manual dishwashing compositions in accord with the invention were prepared. The pH of the compositions was adjusted to be in the range 7.0 to 7.4.

% by weight

	<u>I</u>	<u>II</u>	<u>III</u>	<u>IV</u>	<u>V</u>
LAS	-	-	-	-	10.0
23AE0.8S	10.0	10.0	6.0	5.0	5.0
23AE3S	3.0	7.0	10.0	15.0	-
SS	-	-	4.0	-	-
C12/14 alkyl amine oxide	2.0	1.0	-	1.0	2.0
AEC	-	-	-	5.0	-
C12/14 alkyl di methyl betaine	-	1.0	1.5	2.0	-
C12/14 Ampholak(TM)	-	-	1.5	-	-
CFAA	12.0	-	12.0	11.0	-
APG	-	12.0	-	-	-
C10 Alkyl Ethoxylate (ave. 8)	5.0	5.0	5.0	4.6	5.0
Mg ⁺⁺ ion	-	0.6	0.3	0.3	0.6
Ca ⁺⁺ ion	-	-	0.3	0.15	0.1
Maleic acid	-	-	0.2	0.3	-
Xylanase	0.05	0.1	0.02	0.03	0.04
Protease	0.01	0.02	0.01	0.02	0.03

Water & minors

Up to 100%

Example 3

Granular fabric cleaning compositions in accord with the invention were prepared as follows:

	I	II	III	IV
LAS	22.0	22.0	22.0	22.0
Phosphate	23.0	23.0	23.0	23.0
Carbonate	23.0	23.0	23.0	23.0
Silicate	14.0	14.0	14.0	14.0
Zeolite A	8.2	8.2	8.2	8.2
DETPMP	0.4	0.4	0.4	0.4
Sodium Sulfate	5.5	5.5	5.5	5.5
Protease	0.01	0.02	0.01	0.005
Xylanase	0.04	0.06	0.1	0.04
Lipase	0.005	0.01	-	-
Cellulase	0.001	-	-	0.001
Amylase	0.01	-	0.01	-
Pectinase	0.02	0.02	0.02	-
Water/minors		Up to 100%		

Example 4

Granular fabric cleaning compositions in accord with the invention were prepared as follows:

	I	II	III	IV
LAS	12.0	12.0	12.0	12.0
Zeolite A	26.0	26.0	26.0	26.0
SS	4.0	4.0	4.0	4.0
SAS	5.0	5.0	5.0	5.0
Citrate	5.0	5.0	5.0	5.0
Sodium Sulfate	17.0	17.0	17.0	28.0
Perborate	16.0	16.0	16.0	-
TAED	5.0	5.0	5.0	-
Xylanase	0.20	0.01	0.02	0.08
Protease	0.06	0.03	0.02	0.08
Lipase	0.005	0.01	-	-
Cellulase	0.001	-	-	0.001
Amylase	0.01	-	0.01	-
Pectinase	0.02	0.02	0.02	-
Water/minors		Up to 100%		

Example 5

Granular fabric cleaning compositions in accord with the invention which are especially useful in the laundering of coloured fabrics were prepared as follows :

LAS	11.4	10.7	-
TAS	1.8	2.4	-
TFAA	-	-	4.0
45AS	3.0	3.1	10.0
45E7	4.0	4.0	-
25E3S	-	-	3.0
68E11	1.8	1.8	-
25E5	-	-	8.0
Citrate	14.0	15.0	7.0
Carbonate	-	-	10
Citric acid	3.0	2.5	3.0
Zeolite A	32.5	32.1	25.0
Na-SKS-6	-	-	9.0
MA/AA	5.0	5.0	5.0
DETPMP	1.0	0.2	0.8
Xylanase	0.01	0.05	0.02
Protease	0.02	0.02	0.01
Lipase	0.03	0.04	0.005
Amylase	0.03	0.03	0.005
Pectinase	0.01	-	0.01
Cellulase	0.005	-	0.001
Silicate	2.0	2.5	-
Sulphate	3.5	5.2	3.0
PVP	0.3	0.5	-
Poly (4-vinylpyridine)-N-oxide/copolymer of vinyl-imidazole and vinyl-pyrrolidone	-	-	0.2
Perborate	0.5	1.0	-
Peroxidase	0.01	0.01	-
Phenol sulfonate	0.1	0.2	-
Water/Minors	Up to 100%		

Example 6

Granular fabric cleaning compositions in accord with the invention were prepared as follows:

LAS	6.5	8.0
Sulfate	15.0	18.0
Zeolite A	26.0	22.0
Sodium nitrilotriacetate	5.0	5.0
PVP	0.5	0.7
TAED	3.0	3.0
Boric acid	4.0	-
Perborate	0.5	1.0
Phenol sulphonate	0.1	0.2
Protease	0.06	0.02
Xylanase	0.01	0.02
Silicate	5.0	5.0
Carbonate	15.0	15.0
Peroxidase	0.1	0.1
Lipase	0.01	-
Amylase	0.01	0.01
Pectinase	0.02	-
Cellulase	0.005	0.002
Water/minors	Up to 100%	

Example 7

A compact granular fabric cleaning composition in accord with the invention was prepared as follows:

45AS	8.0
25E3S	2.0
25E5	3.0
25E3	3.0
TFAA	2.5
Zeolite A	17.0
NaSKS-6	12.0
Citric acid	3.0
Carbonate	7.0
MA/AA	5.0
CMC	0.4
Poly (4-vinylpyridine)-N-oxide/ copolymer of vinylimidazole and vinylpyrrolidone	0.2
Xylanase	0.05
Protease	0.05
Lipase	0.005
Cellulase	0.001
Amylase	0.01
TAED	6.0
Percarbonate	22.0
EDDS	0.3
Granular suds suppressor	3.5
water/minors	Up to 100%

Example 8

A granular fabric cleaning compositions in accord with the invention which provide "softening through the wash" capability were prepared as follows:

45AS	-	10.0
LAS	7.6	-
68AS	1.3	-
45E7	4.0	-
25E3	-	5.0
Coco-alkyl-dimethyl hydroxy-ethyl ammonium chloride	1.4	1.0
Citrate	5.0	3.0
Na-SKS-6	-	11.0
Zeolite A	15.0	15.0
MA/AA	4.0	4.0
DETPMP	0.4	0.4
Perborate	15.0	-
Percarbonate	-	15.0
TAED	5.0	5.0
Smectite clay	10.0	10.0
HMWPEO	-	0.1
Protease	0.02	0.01
Lipase	0.02	0.01
Amylase	0.03	0.005
Xylanase	0.03	0.005
Cellulase	0.02	0.001
Pectinase	0.01	0.02
Silicate	3.0	5.0
Carbonate	10.0	10.0
Granular suds suppressor	1.0	4.0
CMC	0.2	0.1
Water/minors	Up to 100%	

Example 9

Heavy duty liquid fabric cleaning compositions suitable for use in the pretreatment of stained fabrics, and for use in a machine laundering method, in accord with the invention were prepared as follows:

	I	II	III	IV	V
24AS	20.0	20.0	20.0	20.0	20.0
SS	5.0	5.0	5.0	5.0	5.0
Citrate	1.0	1.0	1.0	1.0	1.0
12E ₃	13.0	13.0	13.0	13.0	13.0
Monethanolamine	2.5	2.5	2.5	2.5	2.5
Xylanase	0.02	0.01	0.05	0.01	0.03
Protease	0.005	0.03	0.02	0.04	0.01
Lipase	0.002	0.01	0.02	-	0.004
Amylase	0.005	0.005	-	-	0.004
Cellulase	0.04	-	0.01	-	-
Pectinase	0.02	0.02	-	-	-
Water/propylene glycol/ethanol (100:1:1)					

Example 10

Heavy duty liquid fabric cleaning compositions in accord with the invention were prepared as follows:

	I	II	III	IV
LAS acid form	-	-	25.0	-
C ₁₂₋₁₄ alkenyl succinic acid	3.0	8.0	10.0	-
Citric acid	10.0	15.0	2.0	2.0
25AS acid form	8.0	8.0	-	15.0
25AE2S acid form	-	3.0	-	4.0
25AE7	-	8.0	-	6.0
25AE3	8.0	-	-	-
CFAA	-	-	-	6.0
DETPMP	0.2	-	1.0	1.0
Fatty acid	-	-	-	10.0
Oleic acid	1.8	-	1.0	-
Ethanol	4.0	4.0	6.0	2.0
Propanediol	2.0	2.0	6.0	10.0
Xylanase	0.05	0.01	0.005	0.02
Protease	0.02	0.02	0.02	0.01
Amylase	0.005	0.01	0.005	0.01
Cellulase	0.005	-	-	-
Pectinase	0.02	-	-	-
Coco-alkyl dimethyl hydroxy ethyl ammonium chloride	-	-	3.0	-
Smectite clay	-	-	5.0	-
PVP	1.0	2.0	-	-
Perborate	-	1.0	-	-
Phenol sulphonate	-	0.2	-	-
Peroxidase	-	0.01	-	-
NaOH	Up to pH 7.5			
Waters/minors	Up to 100%			

Example 11

The following rinse added fabric softener composition, in accord with the invention, was prepared (parts by weight).

Softener active	24.5
PGMS	2.0
TAE 25	1.5
Xylanase	0.001
Cellulase	0.001
HCL	0.12
Antifoam agent	0.019
Blue dye	80ppm
CaCl ₂	0.35
Perfume	0.90

Example 12aLiquid hard surface cleaning compositions

COMPONENT	A	B
Protease	0.005	0.02
Xylanase	0.01	0.03
Amylase	0.01	0.01
NaC ₁₂ Alkyl-benzene sulfonate	1.95	-
NaC ₁₂ alkylsulfate	-	2.20
NaC ₁₂ (ethoxy)*** sulfate	-	2.20
C ₁₂ dimethylamine oxide	-	0.50
Na Cumene sulfonate	1.30	-
Hexyl Carbitol***	6.30	6.30
Water/minors****	Up to 100%	

*** Diethyleneglycol monohexyl ether

**** All formulas adjusted to pH 7

Example 12bSpray compositions for cleaning hard surfaces
and removing household mildew

COMPONENT	A	B
Protease	0.02	0.005
Amylase	0.01	0.01
Xylanase	0.01	0.02
Sodium octyl sulfate	2.00	2.00
Sodium dodecyl sulfate	4.00	4.00
Sodium hydroxide	0.80	0.80
Silicate (Na)	0.04	0.04
Perfume	0.35	0.35
Water/minors	up to 100%	

Example 12cBar fabric cleaning composition

COMPONENT

Protease	0.03
Xylanase	0.05
C ₁₂ -C ₁₆ alkyl sulfate, Na	20.0
C ₁₂ -C ₁₄ N-methyl glucamide	5.0
C ₁₁ -C ₁₃ alkyl benzene sulfonate, Na	10.0
Sodium carbonate	25.0
Sodium pyrophosphate	7.0
Sodium tripolyphosphate	7.0
Zeolite A	5.0
Carboxymethylcellulose	0.2
Polyacrylate (MW 1400)	0.2
Coconut monethanolamide	5.0
Brightener, perfume	0.2
CaSO ₄	1.0
MgSO ₄	1.0

Filler*

Balance to 100%

* Can be selected from convenient materials such as CaCO₃, talc, clay (Kaolinite, Smectite), silicates and the like.

Example 12dTwo-layer effervescent denture cleansing tablet

COMPONENT

Acidic Layer

Protease	0.1
Xylanase	0.05
Tartaric acid	24.0
Sodium carbonate	4.0
Sulphamic acid	10.0
PEG 20,000	4.0
Sodium bicarbonate	24.5
Potassium persulfate	15.0
Sodium acid pyrophosphate	7.0
Pyrogenic silica	2.0
Tetracetylene diamine	7.0
Ricin-oleylsulfosuccinate	0.5
Flavor	1.0

Alkaline layer

Sodium perborate monohydrate	32.0
Sodium bicarbonate	19.0
EDTA	3.0
Sodium tripolyphosphate	12.0
PEG 20,000	2.0
Potassium persulfate	26.0
Sodium carbonate	2.0
Pyrogenic silica	2.0
Dye/flavor	2.0

Example 13

Fabric swatches (cotton) were stained with banana, pomme granate, grass, tea, clay, tomato puree. After ageing (at least overnight), the stains were split into two equal parts (A & B). Washing of the stains was carried out in a Miele washing machine (cold cycle). One half of the stains were washed with 100g of detergents as given under A (i.e. without xylanase), whereas the second half of the stains were washed with composition B, containing xylanase.

Compositions (all % by weight)

Ingredient	A	B
LAS	6.0	6.0
TAS	2.0	2.0
45E7	4.0	4.0
Zeolite A	16.0	16.0
Citrate	6.0	6.0
Carbonate	15.0	15.0
Silicate	4.0	4.0
MA/AA	4.0	4.0
PB4	20.0	20.0
TAED	5.0	5.0
Granular suds suppressor	2.0	2.0
Protease	0.005	0.005
Lipase	0.005	0.005
Cellulase	0.0005	0.0005
Xylanase	None	60EXU/L wash solution
Water/minors	Up to 100%	

After the wash cycle the swatches were dried and visually graded for stain removal by expert judges using Panel Score Unit (PSU) scale (Scheffe scale).

One xylanase unit (EXU) is defined as the amount of enzyme which, in standard conditions (pH 9.0, 50°C, 30 minutes incubation) degrades xylan.

TABLE I

<u>Stain</u>	<u>PSU (panel score unit) vs NONE xylanase product</u>	
	(B)	(A)
Banana	+ 1.50 (s)	0
Pomme granate	+ 1.13 (s)	0
Tea	+ 1.37 (s)	0
Clay	+ 1.13 (s)	0
Tomato puree	+ 1.00 (s)	0
Grass	+ 0.63 (s)	0

(s) = statistically different

WHAT IS CLAIMED IS:

1. Cleaning composition comprising an enzyme having xylanolytic activity wherein the enzyme is present at a level of from 0.0001% to 2% active enzyme by weight of the composition.
2. Cleaning composition according to claim 1 wherein the enzyme is xylanase.
3. Cleaning composition according to claims 1 or 2 wherein the composition is a detergent composition.
4. Cleaning composition according to claim 3 wherein the detergent composition comprises one or more components selected from nonionic, anionic, cationic, zwitterionic, ampholytic or amphoteric surfactants, bleaching agents, additional enzymes, sud suppressors, dispersants, soil suspension and anti-redeposition agents, smectite clays, builder components and the like.
5. Cleaning composition according to claim 3 or 4 wherein the detergent composition is a granular detergent composition containing no more than 15% by weight of inorganic filler salt.
6. Cleaning composition according to claim 3 wherein the detergent composition is a heavy duty liquid composition.
7. Cleaning composition according to claim 3 wherein the detergent composition is a dishwashing composition.
8. Cleaning composition according to claim 4 wherein the enzymes are selected from the group of cellulases, hemicellulases, peroxidases, proteases, glucoamylases, amylases, lipases, cutinases, pectinases, reductases, oxidases, phenoloxidases, lipoxygenases, ligninases,

pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases or mixtures thereof.

9. Cleaning composition comprising a cocktail of enzymes selected from protease, lipase, cutinase, amylase and/or cellulase in conjunction with one or more plant cell wall degrading enzymes.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/12490

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :C11D 3/386; C12N 9/24, 9/26

US CL :252/174.12, 174.25, 95, 99, DIG. 12;

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 252/174.12, 174.25, 95, 99, DIG. 12;

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

HACHK'S CHEMICAL DICTIONARY

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

search terms : xylanase, xylanolytic, detergent or clean?, laundry, dishwasher?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, A, 92/19726 (SVEDSEN et al) 12 November 1992, see	1-9
---	page 4, line 3, page 10, line 6, and page 17, lines 4-30,	-----
Y	col.18, lines 1-17.	1-9
A	WO, A, 9/01532 (OUTTRUP et al) 20 January 1994, see	1
	col. 6, lines 5-10, col. 8, lines 20-21	
Y	US, A, 5,356,800 (JAQUESS) 18 October 1994, see col. 2,	1-9
	lines 35-54, col. 9, lines 4-13	
A	US, A, 5,290,474 (CLARKSON et al) 01 March 1994, see	1
	col. 2, lines 29-35, col. 10, lines 15 and 60-65, col. 14, line	
	68, and col. 16, line 59-60.	

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

20 NOVEMBER 1995

Date of mailing of the international search report

17 JAN 1996

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer:

PAUL LIEBERMAN

Telephone No. (703) 308-0661

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/12490

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	US, A, 5,434,072 (BOWER) 18 July 1995, see col. 1, lines 60-65, col. 3, lines 5-7, vol. 11, lines 1-20	1-9

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/12490**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 5
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.